

# Demographic and Behavioral Correlates of Cybersickness: A Large Lab-in-the-Field Study of 837 Participants

Tiffany Luong\* Adéla Plechata† Max Möbus\* Michael Atchapero† Robert Böhm†‡§  
Guido Makransky† Christian Holz\*

\*Department of Computer Science, ETH Zürich, Zürich, Switzerland

†Department of Psychology, University of Copenhagen, Copenhagen, Denmark

‡Department of Occupational, Economic, and Social Psychology, University of Vienna, Vienna, Austria

§Copenhagen Center for Social Data Science (SODAS), University of Copenhagen, Copenhagen, Denmark



Figure 1: Our study apparatus was a virtual reality game that informed 837 participants about vaccination benefits. Participants had to navigate a ballroom and avoid contact with wedding guests. After each experience, they reported their cybersickness levels.

## ABSTRACT

Cybersickness has been one of the main impediments to the widespread adoption of Virtual Reality for decades. It has been argued that several factors can influence the occurrence of cybersickness, such as technical factors, interaction design, but also users’ demographics and their perceived presence. Yet, previous studies had comparably small sample sizes and demographically homogeneous samples; comparisons across studies (e.g., regarding demographic factors) are challenging due to the large variation in the studied virtual environments. In this paper, we address these limitations and report the results of a lab-in-the-field experiment on cybersickness with a large and heterogeneous sample of  $N = 837$  participants who navigated and interacted inside a virtual environment (ages 18–80,  $M = 29.34$ ,  $SD = 9.50$ , 431 males, 400 females, 6 non-binaries and other). We found that female participants and participants with lower VR experience were more susceptible to experiencing higher levels of cybersickness. Participants’ cybersickness levels increased with the time spent in VR and with the distance traversed in the virtual world up to a point, above which reported levels declined. We also found a link between higher levels of cybersickness and reduced head motion, as well as between lower levels of cybersickness and more head motion, which led them to explore more of the virtual environment. In contrast to past studies, we did not find any evidence suggesting an effect of age on cybersickness, nor a negative correlation between presence and cybersickness. Based on our results, we derived a model that achieves a mean classification accuracy of 67.1% for two levels of cybersickness using demographic, user experience, and behavioral data in VR.

**Index Terms:** Human-centered computing—Human computer interaction (HCI)—Interaction paradigms—Virtual reality

\*e-mail: firstname.lastname@inf.ethz.ch

†e-mail: firstname.lastname@psy.ku.dk

## 1 INTRODUCTION

Virtual Reality (VR) technologies immerse users in computer-generated 3D worlds, allowing full interaction with the displayed environment, and evoking a sense of presence in the virtual space [63]. This sense of presence opens up the high potential for improvement and development in a wide range of application fields, from training [21] and therapy [5, 62] to social networking [51] and entertainment [2], and can even lead to behavioral changes [46, 73].

One of the main challenges holding VR back from wider adoption is “cybersickness,” the bodily discomfort associated with exposure to immersive content [75]. Cybersickness can produce symptoms of nausea, dizziness, and headaches [57]. These symptoms mimic those of motion sickness, but cybersickness and motion sickness are considered distinct due to their difference in actual physical motion [57] and in symptom severity [66, 67, 69]. Cybersickness has been a known condition in users of virtual and augmented reality systems for decades and even longer in simulators [57]. It impacts user experience and can break users’ immersion in the virtual environments, but can also lead to injury or decreased capacity [57].

While influences of technological and interaction factors on cybersickness have been extensively studied (see survey articles for detailed analyses [12, 57, 71]), previous studies have placed little focus on demographic factors [41, 58]. The lack of evidence regarding a quantified impact of characteristics such as age, presence in VR, and participants’ behavior on cybersickness in combination with the typically small and demographically homogeneous sample sizes of participants in laboratory studies [75] constitutes a major limitation of a more comprehensive understanding of cybersickness. Indeed, Peck et al. [50] and Himmelsbach et al. [26] summarized that past human-computer interaction studies have mostly involved male, white, educated, industrialized, rich, and democratic participants, which significantly under-represents the broader population.

In this paper, we report the results of a large-scale VR intervention study in a museum with a demographically more heterogeneous sample of  $N = 836$  participants (ages 18–30,  $M = 29.34$ ,  $SD = 9.51$ , non-males = 405). Participants were immersed inside a virtual environment that simulated a large indoor event inside a ballroom

during a wedding ceremony as shown in Figure 1. Participants freely navigated the environment and were instructed to complete a series of checkpoints, which led them through multiple rooms, passing roaming people, and finding specific persons to engage in simulated conversations. After completing the VR experience, participants reported their levels of cybersickness and presence in a questionnaire and commented on their VR experience.

Based on the data we collected throughout the study, we derived a cybersickness generalized additive model (GAM) that allowed us to gain insights into occurrences of cybersickness with respect to its demographic and behavioral correlates. Our analysis shows that female participants experienced significantly more cybersickness than male participants and that cybersickness incidence decreased with VR experience. We also found an interaction effect between gender and body ownership. High cybersickness levels for female participants were associated with low body ownership, and high cybersickness levels for male participants were associated with high body ownership. Cybersickness increased with the total duration of experiencing VR, and we found a non-linear relation between the amount of navigation and cybersickness (i.e., total distance traversed in the virtual environment). Overall, cybersickness increased with the amount of navigation in the virtual environment (VE); past a certain point, however, we found a higher amount of navigation to be associated with lower cybersickness scores. Our findings also suggest that participants who reported higher cybersickness ratings moved their heads less while inside VR. Contrarily to past studies, we did not find an influence of age, nor a significant association between presence and cybersickness. Using the variables that had a significant relation with cybersickness, our GAMs achieved a mean accuracy of 67.1% to classify two levels of cybersickness following a user-independent evaluation approach. We conclude with a discussion of our results and put them into perspective.

## 2 RELATED WORK

### 2.1 Demographic Correlates of Cybersickness

Past works that studied the influence of demographic information on cybersickness are summarized in Table 1. Studying the influence of participants' demographic information on cybersickness is not novel, but reports regarding the influence of factors tend to contradict [71]. For example, almost as many studies found sex or gender to have a significant effect on cybersickness [22, 28, 29, 47, 65, 69] as studies that did not find any significant effect of these factors [9, 11, 23, 35, 37, 47, 77] (see Table 1).

Overall, female participants were found to experience more cybersickness than males [22, 28, 29, 47, 65, 69, 71]. This gender effect is in line with studies outside VR as female participants were found to experience more motion sickness and visually induced motion sickness than males [19, 65]. Various explanations have been suggested to explain gender differences in cybersickness. Some explained this difference with the social desirability response bias; male participants can under-report illness or belittle their symptoms to better fit society's gender role expectation [22]. Others attributed the differences to gender dysmorphisms in a seated posture and postural sway [47] or to the non-fitting of the inter-pupillary distance in VR head-mounted displays (HMDs) [65].

Age was also found to influence cybersickness in past studies [1, 22]. This factor was less studied in past work as recruiting a large sample with senior participants can be more difficult [56]. Overall, past studies reported that young users tend to experience less cybersickness than older ones [1, 22, 35]. These results also corroborate past simulator works in non-immersive settings [49].

Past work agrees that habituation can influence cybersickness symptoms. Howarth and Hodder [27] demonstrated that ten exposures to HMD sessions allow most people to adjust to a system. This habituation phenomenon can also extend to video game experience [56, 76] where lower game experience is usually associated

with increased cybersickness incidence [29, 35, 71, 74, 76] (for null effects, see Hakkinen et al. [22]).

There is also a general consensus regarding the influence of past motion sickness history on the incidence of cybersickness symptoms [56, 65, 69]. Motion sickness history is traditionally assessed using the motion history questionnaire (MHQ) [30] or Golding's motion sickness susceptibility questionnaire [16]. In particular, Stanney et al. [69] found a correlation between cybersickness and the carnival ride motion sickness score item from the MHQ.

In general, past studies tend to hold little statistical power due to the relatively small sample size. There are some notable exceptions, such as Stanney et al. [69]'s study in 2003 that gathered 960 participants in a laboratory setting to study the influence of scene complexity, navigation control, gender, body mass index, and motion sickness history on cybersickness. They found that female participants experienced more severe cybersickness symptoms than males, that body mass index was negatively related to oculomotor symptoms, and that cybersickness symptoms ratings increased with the MHQ susceptibility scores.

### 2.2 Other Correlates of Cybersickness

Kolasinski [36] published a list of over two-dozen factors in relation to illness exhibited in simulators. In this section, we report findings about presence and other objective metrics. We refer readers interested in hardware or interaction attributes to prior surveys and reviews on this topic [12, 36, 57].

Duration of exposure was overall found to be positively related to cybersickness (episodes and symptom severity) [28, 36, 45, 69], with works suggesting limiting the duration of VR exposure [12, 69].

However, the relation between presence and cybersickness is far more unclear. Some work found that presence and cybersickness were positively associated [34, 39], negatively associated [1, 76, 78], and some did not find evidence that they were associated [9, 72]. In a recent review, Weech et al. [75] conclude that past findings balance in favor of a negative relationship between presence and cybersickness. However, they mention that high-powered studies could help reveal the nature of this relationship.

Regarding objective metrics, past work mostly focused on physiological modalities such as cardiac activity [3, 34, 42, 44, 82], epidermal activity [42, 44], brain activity [8, 34, 71], respiratory activity [13] or eye-activity [6, 34]. Postural stability has also been extensively explored in the literature as a cause [9, 22, 47] or consequence of cybersickness [23, 69, 82]. However, these metrics can only be assessed using external hardware or modules embedded into VR HMDs [40], which are not always accessible to the general consumer because of their cost and setup requirements.

Regarding modalities that can easily be assessed with current VR HMDs, behavioral measures such as the amount of head motions or interaction in the VE, which is related to interaction and engagement factors [70], have so far received much less attention as potential correlates of cybersickness. We aim to close this gap.

## 3 METHODS

### 3.1 Context of the Study

We conducted the experiment in the National Museum of Berlin over 16 days, from November 11 to 18, 2021. Displays in the museum advertised the study during that time for recruiting purposes. We also advertised and recruited participants on social media networks one month prior to the start of the experiment. In total, 909 participants volunteered to take part in the study or were actively approached in person as they visited the museum.

We designed the experiment for several independent objectives. The first was an entertaining and informative experience about COVID-19 spread and options for vaccination. The details, methods, and results can be found in a submission to a journal in psychology [52]. The second objective of the experiment, investigated in

Table 1: Summary of our study and past VR studies’ findings on the influence of demographic factors on cybersickness. The sources are ordered from most recent (2022) to most ancient (1996). NA=unknown, y=yes, n=no, (empty cells)=not assessed. MS hist.: motion sickness history. SSQ: simulator sickness questionnaire [31]. FMS: fast motion sickness scale [32]. Cust.:custom scale.

Ref.	Year	N	Age			Non-males ratio	VR hardware	Interaction	Context	Seated	Duration (min)	Cybersickness measure	Influence of				
			Range	M	SD								Gender	Age	Game exp.	MS hist.	
Us	2022	837	18-80	29.34	9.50	49%	HMD	joystick	museum	n	~10	Cust.	y	n	y (VR)		
[76]	2020	42	NA	21.74	3.50	36%	HMD	joystick	lab	y	7	SSQ			y		
		128	8-NA	18.2	13.20	64%	HMD	joystick	museum	y	30	SSQ	n		y		
[65]	2020	46	18-30	NA	NA	50%	HMD	none	lab	NA	20	SSQ	y			y	
		147	18-30	NA	NA	50%	HMD	none	lab	NA	20	SSQ	y				
[29]	2020	57	18-38	21.75	NA	37%	HMD	joystick	lab	y	15-	SSQ	y		y		
[11]	2020	79	18-49	21.84	4.19	52%	HMD	simulator, none	NA	y	15	SSQ	n				
[9]	2020	25	18-47	23.92	5.25	48%	HMD	laser pointing	lab	both	15	SSQ, FMS	n				
[77]	2017	29	NA	19.96	NA	48%	HMD	laser pointing	lab	n	10+	SSQ	n				
[47]	2017	36	NA	20.72	0.85	50%	HMD	head movement	lab	y	15-	SSQ	n				
		36	NA	22.72	3.56	50%	HMD	joystick	lab	NA	NA	SSQ	y				
[56]	2014	20	18-31	NA	NA	15%	HMD	joystick	lab	n	NA	SSQ, Cust.			y	y	
[23]	2007	41	NA	NA	NA	49%	HMD	NA	lab	NA	30, 60	NA	n				
[35]	2006	387	9-60	NA	NA	NA	CAVE	NA	tour	NA	NA	SSQ	n	y	y	y	y
[1]	2005	387	9-60	NA	NA	NA	CAVE	NA	tour	NA	NA	SSQ		y			
[69]	2003	960	15-53	21.03	4.43	41%	HMD	mouse	lab	y	15-60	SSQ	y				y
[22]	2002	60	18-41	26.7	NA	20%	HMD	simulator	lab	y	40	SSQ	y	y	n (HMD)		
[28]	2001	60	18-40	NA	NA	30%	HMD	mouse, treadmill	lab	n	15-21+	SSQ	y				
[68]	1999	34	NA	25.79	4.72	41%	HMD	mouse	lab	y	30	SSQ	y				n
[37]	1998	40	19-46	22.7	4.74	50%	HMD	mouse	lab	y	20	SSQ	n				
[36]	1996	40	19-46	22.7	4.74	50%	HMD	mouse	lab	y	20	SSQ	y	y			

the present submission, was to evaluate the impact of demographic and behavioral factors in VR on participants’ rating of presence and cybersickness following the virtual experience.

### 3.2 Virtual Environments

In the VR application, participants were guided through different virtual environments where they had to perform simple interactions. We designed three different versions of the VR experience. The versions participants experienced differed by the presence of gamified elements and/or empathy content. The different versions did not influence the final cybersickness model, however. Each participant experienced one version of the VR experience.

In all versions of the virtual environment, participants embodied an elderly character (gender-matched) who could not get vaccinated against COVID-19 due to medical reasons and was invited to the wedding of his/her granddaughter. We chose elderly avatars as we also aimed at communicating the impact of vaccination on COVID-19 spread for at-risk individuals [52]. Participants were first exposed to an apartment scene in which they were instructed to wash their hands in front of a mirror where they could see their avatar. Then, they received a wedding invitation from their granddaughter.

In the next scene, participants were free to navigate in the 20 × 20 m virtual ballroom (trajectory was not predefined) as shown in Fig. 1. They were guided through different objectives with narration, including signing the guest book, placing a present on a table, and talking to the bride. Participants completed all objectives while following the instruction to actively avoid the other guests, for whom the trajectories were predetermined. We piloted these paths to ensure they were sufficiently complex not to be predicted by participants. When participants accomplished an objective (by reaching a landmark on the floor or by colliding their virtual hand with a highlighted object), the next objective was indicated through a large textbox, arrows, and highlighted landmark in the virtual environment.

For full details on the implementation of the experiment, including the small variations between the three environments, we refer the reader to in our previous work [52].

### 3.3 Apparatus

The support application was developed using Unity and was presented using Oculus Quest 2 in a standalone setting (72 FPS on average). Participants interacted and navigated in the virtual environment using both Oculus Touch. They could move forward and backward using either the controller’s joystick in the direction of their headset (i.e., steering locomotion). When moving, the speed was set to be constant at .75 m/s (not dependent on the joystick motion amplitude) and piloted before the user study with nine persons several times to adjust the game difficulty. There was no deceleration or acceleration in the motion. Participants could pick and drop objects by colliding their avatars’ hands with the 3D models.

The avatars and newlywed models were designed by a professional artist, and the wedding guests were RocketBox avatars [18]. We used voice recordings in either English or German by professional voice actors/actresses for all instructions and interactions with virtual avatars.

### 3.4 Measures

We considered the following data in the analyses:

**Cybersickness:** We asked participants to report their level of motion sickness after the study using the following question: “How motion sick (nauseous or dizzy) did you feel during the wedding experience?” (Scale from 1–Not at all, to 5–Extremely).

**Demographic:** Before the study, we assessed the participants’ gender (1–male, 2–female, 3–non-binary, 4–other), age, VR experience (“How many times have you used Virtual Reality before?” 1–“Never”, 2–“1–3 times”, 3–“4–10 times”, 4–“11–20 times”, 5–“More than 20 times”), and language preference in the VR application (English, German).

**User experience (UX):** After the study, participants reported their level of presence and social presence (adapted from [43]: “While I was at the wedding, I had a sense of “being there”, “I had a sense that I was interacting with other people at the wedding rather than a computer simulation”, scales from 1 (strongly disagree) to 5 (strongly agree), body ownership (from [17]: “I felt as if the virtual body I saw when I looked down was my body”, scale from 1–strongly disagree, to 7–strongly agree).

**Behavioral:** We assessed the total time spent in VR, the accumulated HMD movement (based on the accumulated difference in HMD *position* in the tracking space), the accumulated HMD angular movement (based on the accumulated difference in HMD *rotation* in the tracking space), and the accumulated distance traversed in the VR environment.

Table 2 shows the key sample characteristics.

Table 2: Studied variables and characteristics of the analyzed study sample. Continuous variables are summarized as  $M$  ( $SD$ ).

Variable	Data ( $N = 837$ )
Age [18–80] (in years)	29.34 (9.50)
% Male	51.5% ( $n = 431$ )
% Female	47.8% ( $n = 400$ )
% Non-binary and other	.7% ( $n = 6$ )
Prior VR experience [1–5] (median)	1–Never
% German speakers	60.1% ( $n = 503$ )
Cybersickness [1–5] (median)	2
Presence [1–5] (median)	4
Social presence [1–5] (median)	3
Body ownership [1–7] (median)	5
Time spent in VR (in s)	701.98 (215.83)
HMD movement (in m)	31.32 (13.13)
HMD angular movement (in °)	9905.10 (4434.16)
Distance in VR (in m)	156.53 (48.01)

### 3.5 Participants and Experimental Procedure

In total, 909 participants took part in the experience. 54 participants were excluded from our final sample because they dropped out of the experience due to factors such as time constraints, incomplete or invalid data, or discomfort.

Out of the resulting 855 participants, we excluded 18 additional participants. We excluded nine participants who accumulated an HMD angular movement of more than  $25000^\circ$ , since these observations were larger than the 3<sup>rd</sup> sample quartile plus 1.5 times the inter-quartile range and were found to have a large influence on the model fit (see supplementary materials). For similar reasons, we excluded nine further participants who spent more than 2250 s in VR as part of the experiment (see supplementary materials). Furthermore, we analyzed the data from the participant who reported “other” for gender with the “non-binary” group.

The final sample consisted of  $N = 837$  participants, whose ages ranged from 18 to 80 years (Table 2 lists the sample characteristics).

Participants could choose to complete the simulation and questionnaires either in English or in German. They first filled out an informed consent form and a pre-treatment questionnaire. Then, they were randomly assigned to one of the three VR conditions and spent on average 11.7 min ( $SD = 3.6$  min) in the VR simulation. After the VR experience, participants filled out a post-treatment questionnaire. They were compensated with a small gift (a cup worth 4.50 EUR) and an additional 5 EUR voucher for the museum gift shop.

The study was pre-registered on November 9, 2021, prior to the data collection. It was approved by the relevant Institutional Review Board, approval number IP-IRB/02092021.

## 4 RESULTS

### 4.1 Data Analysis Methods

In this paper, we analyze which demographic, UX and behavioral factors are significantly related to cybersickness. First, we analyzed the relations between variables using Spearman correlations. Then, we modeled cybersickness using a proportional odds logistic regression (POLR) model, since it was reported as an ordered factor on a scale from 1 to 5. In particular, we modeled cybersickness using a POLR GAM, to account for potentially non-linear effects.

Due to their flexibility, GAMs have been a popular modeling technique for complex, high-dimensional data over the last decades and have been constantly enhanced in software such as R [24, 53, 61, 79, 80]. GAMs are particularly frequently used in the fields of environmental, geological, and ecological science [14, 20, 54].

We constructed the POLR GAM for cybersickness using the “MGCV” library [79] in R [53]. The general model included all of the demographic, behavioral, and UX variables indicated in Section 3.4, as well as the different application versions (see Section 3.2). Variables were selected using a sequential backward elimination. The effects with the largest p-values were removed from the model until all effects were statistically significant at a confidence level of .10. Initially, we fitted all effects of not-categorical variables as non-linear effects. If an effect then showed no evidence of non-linear behavior, e.g., was fitted close to a straight line, we subsequently fitted it as a linear effect. Non-linear effects were fitted as thin-plate regression splines, as generally recommended [79, 81]. We included a further penalty on the null space of the regression splines shrinking effects to zero and effectively removing them from the model in case of low explanatory power. This approach allows for variable selection, reduces overfitting of the non-linear effects, and, thus, generally promotes a more conservative fit. In addition to non-linear effects, we also tested for second-order interaction effects. We fitted the GAM using marginal likelihood as the smoothing selection criterion since this results in p-values with the best behavior compared to alternative smoothing selection criteria according to the authors of the used software package “MGCV” [79]. As proposed, we verified that no pair of regression splines exceeded a concurvity of .50 [25].

Finally, we evaluated the cybersickness classification performances using the variables retained in the final model and leave-2-participants-out cross-validation to further test the robustness of the relation between the selected variables and cybersickness.

### 4.2 Correlation Results

The reported cybersickness score correlated statistically significant with participants’ gender ( $\rho = .28, p < .001$ ), previous VR experience ( $\rho = -.15, p < .001$ ), total duration in VR as part of the experiment ( $\rho = .08, p < .05$ ) and accumulated HMD angular movement ( $\rho = -.08, p < .05$ ). No other correlations were statistically different to 0, at a confidence level lower than .05. A full correlation matrix between all collected variables can be found in Fig. 2.




Figure 2: Correlation matrix of all included variables. The data distributions, detailed correlation coefficients (Spearman’s  $\rho$ ), and p-values can be found in supplementary materials.

### 4.3 Cybersickness Model

In the following sections, we will discuss the variables that were calculated to have a statistically significant effect on cybersickness in the POLR GAM. Following the procedure indicated in Section 4.1, we obtained the following model:

$$\begin{aligned} \text{cybersickness} &= g(\eta), \\ \text{with } \eta_i &= \text{gender}_i + \text{VR\_experience}_i \\ &\quad + \text{body\_ownership}_i \\ &\quad + \text{body\_ownership}_i : \text{gender}_i \\ &\quad + \text{time\_in\_VR}_i + f_1(\text{hmd\_rotation}_i) \\ &\quad + f_2(\text{distance\_in\_VR}_i) \text{ for } i \in 1, \dots, n, \end{aligned} \quad (1)$$

where  $f_j$  are smooth functions representing the effects of  $\text{hmd\_rotation}$  and  $\text{distance\_in\_VR}$  for  $j = 1$  and  $j = 2$ , respectively.

If  $\tau_2, \tau_3, \tau_4 \in \mathbf{R}$  are three distinct points on the real line,  $g$  takes the following form for an ordinal response with 5 levels  $(1, \dots, 5)$ :

$$g(\eta) = \begin{cases} 1, & \text{if } \eta < -1 \\ 2, & \text{if } -1 \leq \eta < \tau_2 \\ 3, & \text{if } \tau_2 \leq \eta < \tau_3 \\ 4, & \text{if } \tau_3 \leq \eta < \tau_4 \\ 5, & \text{otherwise} \end{cases} \quad (2)$$

For our final model (POLR GAM):

$$\tau_2 = 0.77; \quad \tau_3 = 2.04; \quad \tau_4 = 4.14 \quad (3)$$

The partial dependence plots for each variable can be found in Fig. 3.

#### 4.3.1 Demographic and UX Factors

We found a statistically significant effect of gender on the reported cybersickness score (see Fig. 3). Female participants were calculated to have increased chances of experiencing cybersickness compared to male participants ( $p < .0001$ ,  $\chi_2^2 = 24.56$ ), matching previous observations of Spearman correlations. While participants self-identifying as non-binary or other were also calculated to experience cybersickness more easily than male participants, this difference was not statistically significant. We found previous experience in VR to significantly decrease the chances of experiencing cybersickness ( $p < .0001$ ,  $\chi_1^2 = 23.97$ ; see Fig. 3)—again matching previously observed Spearman correlations. For female participants, our findings also showed that decreases in experienced body ownership during the experiment significantly increased the chances of observing high levels of cybersickness ( $p < .01$ ; see Fig. 3). For male participants, however, the trend was reversed as increased body ownership tended to increase the expected cybersickness measure ( $p = .10$ ; see Fig. 3). For participants identifying as non-binary or other, the effect of body ownership was not statistically significant ( $p = .92$ ).

We did not find any significant effect of age nor presence on cybersickness (see Fig. 4).

#### 4.3.2 Behavioral Variables

Furthermore, we found multiple statistically significant effects connected to the behavior of participants during the experiment: total time spent in VR, total distance traversed in VR, and accumulated HMD angular movement. Increases in total time spent in VR increased the chances of experiencing cybersickness linearly ( $p = .01$ ,  $\chi_1^2 = 6.61$ ). Increases in accumulated HMD angular movement decreased the expected cybersickness measure non-linearly ( $p < .01$ ,  $s(edf) = 1.80$ ,  $\chi_1^2 = 9.34$ ) as can be seen in Fig. 3. The fitted effect decreased steepest in between  $\approx 10,000^\circ$ – $15,000^\circ$ . Total distanced traversed in VR also showed evidence of a non-linear effect ( $p < .01$ ,  $s(edf) = 2.14$ ,  $\chi_1^2 = 9.43$ ). We observed an effect similar to a negative quadratic parabola: traversing only short distances in VR (less

than 100 m) was calculated to decrease the expected cybersickness measure, as well as traversing long distances (more than 300 m). We found traversing distances around 200 m to maximize the chances of observing cybersickness.

We did not find that other behavioral variables were significantly related to cybersickness.

### 4.4 Classification Performances

We explored whether the variables retained in the final GAM (Equation 1) could be used to classify cybersickness by performing a leave-2-participants-out cross-validation (1000 iterations, balanced test sets). We compared the performances of POLR GAMs, support vector machines (SVM), and random forests (RF; 500 estimators and maximum depth of 5). The evaluation was performed for 2 classes of cybersickness levels ([1,2] vs. [3,4,5]), and the training sets were balanced using Synthetic Minority Oversampling Technique (SMOTE) [7].

Table 3 summarizes our results.

Table 3: Cybersickness classification performances.

N Classes		2	
Cybersickness ratings		[1,2] vs. [3,4,5]	
Metric		Accuracy	micro F1 score
Algorithm	POLR	67.1%	66.6%
	SVM	64.5%	64.5%
	RF	59.3%	59.2%

The results show that the POLR GAMs overall outperformed the SVM and RF models with a mean accuracy of 67.1%.

## 5 DISCUSSION

In this paper, we conducted a study and assessed participants' cybersickness level, as well as their demographic information and behavioral data. The richness of this analysis stems from the exceptionally large ( $N = 837$ ) and diverse ( $SD_{age} = 9.50$ ,  $\text{non-males}_{ratio} = 48.5\%$ ) sample of participants who were recruited in the wild in a museum, outside laboratory settings. Below we summarize the results on factors being associated with participants' self-reported levels of cybersickness; we put them into perspectives of the previous literature.

### 5.1 Demographic and UX Correlates

Female participants reported significantly higher levels of cybersickness than male participants, which corroborates most of the past findings [22, 28, 29, 47, 65, 69]. Stanney et al. [65] explained the gender effect with the non-fit of the IPD and Munafo et al. [47] with participants' postural sway, which we did not control in this study. The HMD was calibrated vertically using the Oculus Quest 2 procedure, but the distance between the lenses was not adjusted between participants. Another potential explanation is social desirability biases linked to society's gender role expectation [10, 15, 22]. Out of the 431 male participants, *none* reported the maximum level of cybersickness, i.e., 5. Given the exceptionally large and diverse sample, these findings still strengthen past results that found that female users are more susceptible to experiencing cybersickness than male users, also using the most widespread VR HMD on the market in 2022.

In contrast to past studies [1, 22, 35], we did not observe a relation between age and cybersickness (see Fig. 2 and the supplementary materials). Some of these previous studies were brief reports and did not report important details (e.g., age mean and SD, test statistics) or were limited by low sample size. Taken together, previous estimates are likely to be less accurate, over- or underestimating the true effect.






Figure 3: Partial dependence plots of the effects of the dependent variables on cybersickness in the final GAM model (equation 1). The y-axis corresponds to the change in the linear predictor  $\eta$  (equation 1), which is transformed into the average cybersickness level using function  $g$  (equation 2) with cut-off values from equation 3.  $s(edf)$  reflects the effective degrees of freedom of the fitted smoothing splines for non-linear effects relating to the complexity of the fitted non-linear effect. Linear effects require one degree of freedom, indicated by  $p(1)$ . Similarly,  $c(edf)$  refers to the degrees of freedom used by categorical variables. Dotted lines indicate 95% confidence intervals of the fits. Only significant effects ( $p < .05$ ) are colored. Stacked bar charts for gender, prior VR experience, and body ownership can be found in supplementary materials.

Our results show that an increase in past VR exposures decreased the level of cybersickness reported. Few studies assessed participants’ VR experience (they mostly measured game experience [29, 35, 76]), and the only work that assessed it did not find a significant effect of VR experience on cybersickness [22]. Our results can mainly be explained by the phenomena of habituation [27, 59]. Additionally, it is more likely that people who experienced cybersickness in the past will less self-select into an experience using a VR HMD again.

We did not find a correlation between presence (environment and social) and cybersickness (see Fig. 2 and 3). There have been debates about such a relation in the past [9]. A recent review settled for a negative correlation between presence and cybersickness but the authors highlighted the lack of “high-powered studies” [75]. The null correlation might be due to a complicated causal-inference relation on/of cybersickness for different participants. Weech et al. [75] mentioned that a positive relationship could occur because of the mutual impact of immersiveness—the sensory “submersion” experienced by a user [4] or vection—the illusion of self-motion [60] on presence and on cybersickness. These factors are sometimes considered as preconditions for users to experience high levels of presence [45, 60, 64] or high levels of cybersickness (through sensory conflicts) [38, 55]. In contrast, a negative relation between cybersickness and presence can be explained by a top-down relation, i.e., cybersickness could have played a distracting role in the VR experience, suppressing the attention required to achieve high levels of presence [48, 60, 72, 75].

The results also show an interaction effect between gender and body ownership on cybersickness. Female participants with *lower* body ownership experienced significantly higher degrees of cybersickness, while male participants with *higher* body ownership tended

to experience higher degrees of cybersickness. A similar interpretation as the one for the positive or negative relation between presence and cybersickness can be given for this interaction effect. On the one hand, the positive association between body ownership and cybersickness for male participants could originate from an external factor such as vection or immersiveness. The illusion of self-motion requires the sensorimotor control to be convinced that the visual motion is in line with one’s own body motion [60]. In that sense, body ownership is even more concerned with vection than the overall sense of presence. On the other hand, cybersickness might have prevented female participants to experience the illusion of body ownership. A past work found that female participants experienced a higher onset of cybersickness symptoms than males [23]. This might also have been the case in our study, which could have impacted females’ user experience at an earlier point. Further work could focus on studying this potential difference in cybersickness and body ownership outcome, based on when cybersickness occurred.

## 5.2 Behavioral Correlates

We found a positive association between the time spent in VR and cybersickness, corroborating past results [28, 36, 45, 69].

Interestingly, we found a negative non-linear association between accumulated HMD angular movement and cybersickness ratings. This effect could seem counter-intuitive from a bottom-up point of view as one could expect a higher amount of interaction and motion to increase the level of cybersickness. The most plausible explanation is that the relation originates from cybersickness (i.e., top-down effect). Participants who experienced cybersickness might have decreased their head movement in VR to avoid worsening their sickness symptoms while a higher amount of HMD rotation




Figure 4: Non-significant association (left) between age and cybersickness ratings, (middle) between presence in an environment and cybersickness, and (right) between social presence and cybersickness.

translated to a lower chance of cybersickness occurrence and higher confidence in interacting in the VR environment (looking around). There are parallels to draw with past results. Indeed, Steinicke et al. [70] found that looking in the direction of the walk indicated that users felt insecure, whereas a free-look-around revealed that users felt safe. Further studies could focus on these interpretations.

Furthermore, the total distance traversed in VR is also significantly associated with the level of cybersickness. More specifically, the level of cybersickness is positively related to the total distance traveled in VR up to a point ( $\sim 200$  m), above which the reported levels declined. Some interpretations: (1) bottom-up in the first part, the cause of this relationship might originate from the interaction, where the more users perform navigation interaction, the higher is their level of cybersickness; (2) top-down in the second part, non-sick participants were more at-ease with interacting in VR and felt free to explore the VR environment to their heart content, similarly to the HMD angular movement results. These effects require further investigation.

### 5.3 Limitations and Future Work

Our results revealed interesting insights regarding the incidence of cybersickness in—what could be considered a regular VR application showcase for passerby visitors in a museum. We discovered significant relations between behavioral metrics and cybersickness, which pave the way for further research. In particular, our results suggest that while longer VR exposure augments the risk to experience high levels of cybersickness, a large amount of HMD rotation and exploration in VR translates to low levels of cybersickness. Consequently, an interesting question to investigate would be: does cybersickness decrease users’ head rotation? Furthermore, we only assessed sum metrics for behavioral measures. The joint effect of HMD rotation and users’ navigation was not sampled and could be further investigated to better understand sensory conflicts in VR in the future. Overall, a big question stems from these results: how can we disentangle the complex relation betweenvection, presence, body ownership, users’ behavior, and cybersickness? Those are still unanswered questions that would require additional experiments.

An important limit in our method is the assessment of cybersickness. We used a non-standardized single state question to assess cybersickness, as opposed to the widely spread simulator sickness questionnaire (SSQ) [31]. We did not use the SSQ as the user study was constrained in time (i.e.,  $< 20$  min per participant) as part of an exhibition in a museum. In the question, we specified nausea and dizziness symptoms as those weight the most with general discomfort in the SSQ final score [31]. Still, it is unclear from our assessment to what extent participants felt either or both and some symptoms such as oculomotor ones were not assessed. A shorter standardized scale for VR cybersickness would benefit future large-scale VR experiments and longer questionnaires such as the SSQ or VRSQ [33] could improve the diagnosticity for cybersickness symptoms.

We reported the demographic, UX, and behavioral factors that

were significantly related to cybersickness in our study. Using POLR GAMs and machine learning approaches, we reached a mean classification accuracy of 67.1% for two levels of cybersickness, highlighting the potential of the identified variables to predict cybersickness.

Future work could consider several lines for improvement. Especially, physiological modalities have shown promises to detect cybersickness [8, 13, 34, 44, 71]. In our study, we only considered data that could be obtained from the standalone Oculus Quest 2 out of the box. Future work could attempt to establish a large dataset on cybersickness studies in VR with demographic, behavioral, and physiological data to better understand the individual variability of cybersickness and to detect cybersickness in real-time.

Overall, our results are reassuring as they support past findings with a smaller sample size regarding the effect of gender, VR experience, and time in VR. However, they also contradict past work on the relation between cybersickness and age and presence. Therefore, we argue that large sample sizes should be employed whenever possible in future cybersickness studies, especially when demographics are studied because of the large individual variabilities.

## 6 CONCLUSION

In this paper, we conducted a large lab-in-the-field study in a museum with  $N = 837$  participants and analyzed the relation between cybersickness and demographic, user experience, and behavioral measures.

Our results show that female participants and participants with low VR experience experienced higher levels of cybersickness. High cybersickness level for female participants was associated with low body ownership, and high cybersickness level for male participants was associated with high body ownership. We confirm that a longer time in virtual reality (VR) and an increased amount of navigation in VR—until a certain point, increase the risk of experiencing higher levels of cybersickness. Our results also show a positive association between low cybersickness level and increased amount of head rotation, as well as between low cybersickness level and increased distance traversed in the immersive virtual environment. Unlike past work, we did not find an effect of age on cybersickness, nor a significant association between presence and cybersickness. Using the variables that had a significant relation with cybersickness, our models achieved a mean classification accuracy of 67.1% for two levels of cybersickness on unseen participants.

Taken together, our results suggest that female participants and participants with low VR experience are more susceptible to experiencing cybersickness. Some factors such as the time spent in VR and the quantity of navigation in VR can also increase the risk of cybersickness incidence. Our findings suggest that participants who experienced low levels of cybersickness moved their heads and explored the immersive virtual environment more. Those results pave the way for future experiments aiming at understanding the relation between cybersickness, users’ behaviors,vection, presence, and body ownership to better predict cybersickness occurrences.

## ACKNOWLEDGMENTS

This research was funded by the European Institute of Innovation and Technology (EIT Health, Grant No. 210836). We would also like to warmly thank the National Museum of Berlin and the students from the University of Copenhagen who helped conducting this study.

## REFERENCES

- [1] Laura L Arns and Melinda M Cerney. The relationship between age and incidence of cybersickness among immersive environment users. In *IEEE Proceedings. VR 2005. Virtual Reality, 2005.*, pages 267–268. IEEE, 2005.
- [2] Ruth Aylett and Sandy Louchart. Towards a narrative theory of virtual reality. *Virtual Reality*, 7(1):2–9, 2003.
- [3] RJV Bertin, C Collet, S Espié, and W Graf. Objective measurement of simulator sickness and the role of visual-vestibular conflict situations. In *Driving Simulation Conference North America*, pages 280–293. University of Central Florida Orlando, USA, 2005.
- [4] Frank Biocca and Ben Delaney. Immersive virtual reality technology. *Communication in the age of virtual reality*, 15(32):10–5555, 1995.
- [5] Cristina Botella, Javier Fernández-Álvarez, Verónica Guillén, Azucena García-Palacios, and Rosa Baños. Recent progress in virtual reality exposure therapy for phobias: a systematic review. *Current psychiatry reports*, 19(7):1–13, 2017.
- [6] Eunhee Chang, Hyun Taek Kim, and Byounghyun Yoo. Predicting cybersickness based on user’s gaze behaviors in hmd-based virtual reality. *Journal of Computational Design and Engineering*, 8(2):728–739, 2021.
- [7] Nitesh V Chawla, Kevin W Bowyer, Lawrence O Hall, and W Philip Kegelmeyer. Smote: synthetic minority over-sampling technique. *Journal of artificial intelligence research*, 16:321–357, 2002.
- [8] Yu-Chieh Chen, Jeng-Ren Duann, Shang-Wen Chuang, Chun-Ling Lin, Li-Wei Ko, Tzyy-Ping Jung, and Chin-Teng Lin. Spatial and temporal eeg dynamics of motion sickness. *NeuroImage*, 49(3):2862–2870, 2010.
- [9] Jeremy Clifton and Stephen Palmisano. Effects of steering locomotion and teleporting on cybersickness and presence in hmd-based virtual reality. *Virtual Reality*, 24(3):453–468, 2020.
- [10] Will H Courtenay. Constructions of masculinity and their influence on men’s well-being: a theory of gender and health. *Social science & medicine*, 50(10):1385–1401, 2000.
- [11] Christopher Curry, Ruixuan Li, Nicolette Peterson, and Thomas A Stoffregen. Cybersickness in virtual reality head-mounted displays: examining the influence of sex differences and vehicle control. *International Journal of Human-Computer Interaction*, 36(12):1161–1167, 2020.
- [12] Simon Davis, Keith Nesbitt, and Eugene Nalivaiko. A systematic review of cybersickness. In *Proceedings of the 2014 conference on interactive entertainment*, pages 1–9, 2014.
- [13] Mark S Dennison, A Zachary Wisti, and Michael D’Zmura. Use of physiological signals to predict cybersickness. *Displays*, 44:42–52, 2016.
- [14] Francesca Dominici, Aidan McDermott, Scott L. Zeger, and Jonathan M. Samet. On the Use of Generalized Additive Models in Time-Series Studies of Air Pollution and Health. *American Journal of Epidemiology*, 156(3):193–203, 08 2002. ISSN 0002-9262.
- [15] Joan Evans, Blye Frank, John L Oliffe, and David Gregory. Health, illness, men and masculinities (himm): a theoretical framework for understanding men and their health. *Journal of Men’s Health*, 8(1):7–15, 2011.
- [16] John F Golding. Motion sickness susceptibility questionnaire revised and its relationship to other forms of sickness. *Brain research bulletin*, 47(5):507–516, 1998.
- [17] Mar Gonzalez-Franco and Tabitha C Peck. Avatar embodiment towards a standardized questionnaire. *Frontiers in Robotics and AI*, 5:74, 2018.
- [18] Mar Gonzalez-Franco, Eyal Ofek, Ye Pan, Angus Antley, Anthony Steed, Bernhard Spanlang, Antonella Maselli, Domna Banakou, Nuria Pelechano, Sergio Orts-Escolano, et al. The rocketbox library and the utility of freely available rigged avatars. *Frontiers in virtual reality*, page 20, 2020.
- [19] David A Graeber and Kay M Stanney. Gender differences in visually induced motion sickness. In *Proceedings of the human factors and ergonomics society annual meeting*, volume 46, pages 2109–2113. SAGE Publications Sage CA: Los Angeles, CA, 2002.
- [20] Antoine Guisan, Thomas C Edwards Jr, and Trevor Hastie. Generalized linear and generalized additive models in studies of species distributions: setting the scene. *Ecological modelling*, 157(2-3):89–100, 2002.
- [21] Kurinchi Selvan Gurusamy, Rajesh Aggarwal, Latha Palanivelu, and Brian R Davidson. Virtual reality training for surgical trainees in laparoscopic surgery. *Cochrane database of systematic reviews*, (1), 2009.
- [22] Jukka Hakkinen, Tero Vuori, and M Paakka. Postural stability and sickness symptoms after hmd use. In *IEEE International Conference on Systems, Man and Cybernetics*, volume 1, pages 147–152. IEEE Yasmine Hammamet, 2002.
- [23] DL Harm, LC Taylor, and JJ Bloomberg. Adaptive changes in sensorimotor coordination and motion sickness following repeated exposures to virtual environments. In *NASA Human Research Program Investigators’ Meeting*, 2007.
- [24] Trevor Hastie. *gam: Generalized Additive Models*, 2022. URL <https://CRAN.R-project.org/package=gam>.
- [25] Shui He. *Generalized additive models for data with concavity: Statistical issues and a novel model fitting approach*. PhD thesis, University of Pittsburgh, 2004.
- [26] Julia Himmelsbach, Stephanie Schwarz, Cornelia Gerdenitsch, Beatrix Wais-Zechmann, Jan Bobeth, and Manfred Tscheligi. Do we care about diversity in human computer interaction: A comprehensive content analysis on diversity dimensions in research. In *Proceedings of the 2019 CHI Conference on Human Factors in Computing Systems*, pages 1–16, 2019.
- [27] Peter A Howarth and Simon G Hodder. Characteristics of habituation to motion in a virtual environment. *Displays*, 29(2):117–123, 2008.
- [28] Beverly K Jaeger and Ronald R Mourant. Comparison of simulator sickness using static and dynamic walking simulators. In *Proceedings of the Human Factors and Ergonomics Society Annual Meeting*, volume 45, pages 1896–1900. SAGE Publications Sage CA: Los Angeles, CA, 2001.
- [29] Angelica Jasper, Nicholas Cone, Chase Meusel, Michael Curtis, Michael C Dorneich, and Stephen B Gilbert. Visually induced motion sickness susceptibility and recovery based on four mitigation techniques. *Frontiers in Virtual Reality*, 1:22, 2020.
- [30] Robert S Kennedy, Jennifer E Fowlkes, Kevin S Berbaum, and Michael G Lilienthal. Use of a motion sickness history questionnaire for prediction of simulator sickness. *Aviation, Space, and Environmental Medicine*, 63(7):588–593, 1992.
- [31] Robert S Kennedy, Norman E Lane, Kevin S Berbaum, and Michael G Lilienthal. Simulator sickness questionnaire: An enhanced method for quantifying simulator sickness. *The international journal of aviation psychology*, 3(3):203–220, 1993.
- [32] Behrang Keshavarz and Heiko Hecht. Validating an efficient method to quantify motion sickness. *Human factors*, 53(4):415–426, 2011.



- [33] Hyun K Kim, Jaehyun Park, Yeongcheol Choi, and Mungyeong Choe. Virtual reality sickness questionnaire (vrsq): Motion sickness measurement index in a virtual reality environment. *Applied ergonomics*, 69:66–73, 2018.
- [34] Young Youn Kim, Hyun Ju Kim, Eun Nam Kim, Hee Dong Ko, and Hyun Taek Kim. Characteristic changes in the physiological components of cybersickness. *Psychophysiology*, 42(5):616–625, 2005.
- [35] Melinda M Knight and Laura L Arns. The relationship among age and other factors on incidence of cybersickness in immersive environment users. In *Proceedings of the 3rd Symposium on Applied Perception in Graphics and Visualization*, pages 162–162, 2006.
- [36] Eugenia M Kolasinski. *Prediction of simulator sickness in a virtual environment*. University of Central Florida, 1996.
- [37] Eugenia M Kolasinski and Richard D Gilson. Simulator sickness and related findings in a virtual environment. In *Proceedings of the human factors and ergonomics society annual meeting*, volume 42, pages 1511–1515. SAGE Publications Sage CA: Los Angeles, CA, 1998.
- [38] JJ-W Lin, Henry Been-Lirn Duh, Donald E Parker, Habib Abi-Rached, and Thomas A Furness. Effects of field of view on presence, enjoyment, memory, and simulator sickness in a virtual environment. In *Proceedings IEEE Virtual Reality 2002*, pages 164–171. IEEE, 2002.
- [39] Cheng-Li Liu and Shiao-Tsyur Uang. Effects of presence on causing cybersickness in the elderly within a 3d virtual store. In *International conference on human-computer interaction*, pages 490–499. Springer, 2011.
- [40] Tiffany Luong, Anatole Lecuyer, Nicolas Martin, and Ferran Argelaguet. A survey on affective and cognitive vr. *IEEE Transactions on Visualization and Computer Graphics*, 2021.
- [41] Cayley MacArthur, Arielle Grinberg, Daniel Harley, and Mark Hancock. You're making me sick: A systematic review of how virtual reality research considers gender & cybersickness. In *Proceedings of the 2021 CHI Conference on Human Factors in Computing Systems*, pages 1–15, 2021.
- [42] Takuro Magaki and Michael Vallance. Developing an accessible evaluation method of vr cybersickness. In *2019 IEEE Conference on Virtual Reality and 3D User Interfaces (VR)*, pages 1072–1073. IEEE, 2019.
- [43] Guido Makransky, Lau Lilleholt, and Anders Aaby. Development and validation of the multimodal presence scale for virtual reality environments: A confirmatory factor analysis and item response theory approach. *Computers in Human Behavior*, 72:276–285, 2017.
- [44] Nicolas Martin, Nicolas Mathieu, Nico Pallamin, Martin Ragot, and Jean-Marc Diverrez. Virtual reality sickness detection: an approach based on physiological signals and machine learning. In *2020 IEEE International Symposium on Mixed and Augmented Reality (ISMAR)*, pages 387–399. IEEE, 2020.
- [45] Michael E McCauley and Thomas J Sharkey. Cybersickness: Perception of self-motion in virtual environments. *Presence: Teleoperators & Virtual Environments*, 1(3):311–318, 1992.
- [46] Aske Mottelson, Clara Vandeweerd, Michael Atchapero, Tiffany Luong, Christian Holz, Robert Böhm, and Guido Makransky. A self-administered virtual reality intervention increases covid-19 vaccination intention. *Vaccine*, 39(46):6746–6753, 2021.
- [47] Justin Munafo, Meg Diedrick, and Thomas A Stoffregen. The virtual reality head-mounted display oculus rift induces motion sickness and is sexist in its effects. *Experimental brain research*, 235(3):889–901, 2017.
- [48] Sarah Nichols, Clovissa Haldane, and John R Wilson. Measurement of presence and its consequences in virtual environments. *International Journal of Human-Computer Studies*, 52(3):471–491, 2000.
- [49] George D Park, R Wade Allen, Dary Fiorentino, Theodore J Rosenthal, and Marcia L Cook. Simulator sickness scores according to symptom susceptibility, age, and gender for an older driver assessment study. In *Proceedings of the human factors and ergonomics society annual meeting*, volume 50, pages 2702–2706. SAGE Publications Sage CA: Los Angeles, CA, 2006.
- [50] Tabitha C Peck, Kyla A McMullen, and John Quarles. Divrsify: Break the cycle and develop vr for everyone. *IEEE Computer Graphics and Applications*, 41(6):133–142, 2021.
- [51] Tekla S Perry. Virtual reality goes social. *IEEE Spectrum*, 53(1):56–57, 2015.
- [52] Adéla Plechatá, Clara Vandeweerd, Michael Atchapero, Tiffany Luong, Christian Holz, Cornelia Betsch, Bonnie Dietermann, Yori Schultka, Robert Böhm, and Guido Makransky. Experiencing herd immunity in virtual reality increases covid-19 vaccination intention: Evidence from a large-scale field intervention study, May 2022. URL [psyarxiv.com/68wqn](https://psyarxiv.com/68wqn).
- [53] R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2021. URL <https://www.R-project.org/>.
- [54] Khaiwal Ravindra, Preeti Rattan, Suman Mor, and Ashutosh Nath Aggarwal. Generalized additive models: Building evidence of air pollution, climate change and human health. *Environment International*, 132:104987, 2019. ISSN 0160-4120.
- [55] James T Reason and Joseph John Brand. *Motion sickness*. Academic press, 1975.
- [56] Lisa Rebenitsch and Charles Owen. Individual variation in susceptibility to cybersickness. In *Proceedings of the 27th annual ACM symposium on User interface software and technology*, pages 309–317, 2014.
- [57] Lisa Rebenitsch and Charles Owen. Review on cybersickness in applications and visual displays. *Virtual Reality*, 20(2):101–125, 2016.
- [58] Lisa Rebenitsch and Charles Owen. Estimating cybersickness from virtual reality applications. *Virtual Reality*, 25(1):165–174, 2021.
- [59] Clare Regan. An investigation into nausea and other side-effects of head-coupled immersive virtual reality. *Virtual Reality*, 1(1):17–31, 1995.
- [60] Bernhard E Riecke. Compelling self-motion through virtual environments without actual self-motion: using self-motion illusions (“vection”) to improve user experience in vr. *Virtual reality*, 8(1):149–178, 2011.
- [61] R. A. Rigby and D. M. Stasinopoulos. Generalized additive models for location, scale and shape,(with discussion). *Applied Statistics*, 54:507–554, 2005.
- [62] Giuseppe Riva. Virtual reality in psychotherapy. *Cyberpsychology & behavior*, 8(3):220–230, 2005.
- [63] Mel Slater, Martin Usoh, and Anthony Steed. Depth of presence in virtual environments. *Presence: Teleoperators & Virtual Environments*, 3(2):130–144, 1994.
- [64] Mel Slater, Vasilis Linakis, Martin Usoh, and Rob Kooper. Immersion, presence and performance in virtual environments: An experiment with tri-dimensional chess. In *Proceedings of the ACM symposium on virtual reality software and technology*, pages 163–172, 1996.
- [65] Kay Stanney, Cali Fidopiastis, and Linda Foster. Virtual reality is sexist: but it does not have to be. *Frontiers in Robotics and AI*, 7:4, 2020.
- [66] Kay M Stanney and Robert S Kennedy. The psychometrics of cybersickness. *Communications of the ACM*, 40(8):66–68, 1997.

- [67] Kay M Stanney, Robert S Kennedy, and Julie M Drexler. Cybersickness is not simulator sickness. In *Proceedings of the Human Factors and Ergonomics Society annual meeting*, volume 41, pages 1138–1142. SAGE Publications Sage CA: Los Angeles, CA, 1997.
- [68] Kay M Stanney, Robert S Kennedy, Julie M Drexler, and Deborah L Harm. Motion sickness and proprioceptive aftereffects following virtual environment exposure. *Applied ergonomics*, 30(1):27–38, 1999.
- [69] Kay M Stanney, Kelly S Hale, Isabelina Nahmens, and Robert S Kennedy. What to expect from immersive virtual environment exposure: Influences of gender, body mass index, and past experience. *Human factors*, 45(3):504–520, 2003.
- [70] Frank Steinicke, Gerd Bruder, Klaus Hinrichs, Anthony Steed, and Alexander L Gerlach. Does a gradual transition to the virtual world increase presence? In *2009 IEEE Virtual Reality Conference*, pages 203–210. IEEE, 2009.
- [71] Nana Tian, Phil Lopes, and Ronan Boulic. A review of cybersickness in head-mounted displays: raising attention to individual susceptibility. *Virtual Reality*, pages 1–33, 2022.
- [72] Martin Usoh, Kevin Arthur, Mary C Whitton, Rui Bastos, Anthony Steed, Mel Slater, and Frederick P Brooks Jr. Walking, walking-in-place, flying, in virtual environments. In *Proceedings of the 26th annual conference on Computer graphics and interactive techniques*, pages 359–364, 1999.
- [73] Clara Vandeweerd, Tiffany Luong, Michael Atchapero, Aske Mottelson, Christian Holz, Guido Makransky, and Robert Böhm. Virtual reality reduces covid-19 vaccine hesitancy in the wild: a randomized trial. *Scientific Reports*, 12(1):1–7, 2022.
- [74] Yuyang Wang, Jean-Rémy Chardonnet, Frédéric Merienne, and Jivka Ovtcharova. Using fuzzy logic to involve individual differences for predicting cybersickness during vr navigation. In *2021 IEEE Virtual Reality and 3D User Interfaces (VR)*, pages 373–381. IEEE, 2021.
- [75] Séamas Weech, Sophie Kenny, and Michael Barnett-Cowan. Presence and cybersickness in virtual reality are negatively related: a review. *Frontiers in psychology*, 10:158, 2019.
- [76] Séamas Weech, Sophie Kenny, Markus Lenizky, and Michael Barnett-Cowan. Narrative and gaming experience interact to affect presence and cybersickness in virtual reality. *International Journal of Human-Computer Studies*, 138:102398, 2020.
- [77] Michael L Wilson and Amelia J Kinsela. Absence of gender differences in actual induced hmd motion sickness vs. pretrial susceptibility ratings. In *Proceedings of the Human Factors and Ergonomics Society Annual Meeting*, volume 61, pages 1313–1316. SAGE Publications Sage CA: Los Angeles, CA, 2017.
- [78] Bob G Witmer, John H Bailey, Bruce W Knerr, and Kimberly C Parsons. Virtual spaces and real world places: Transfer of route knowledge. *International journal of human-computer studies*, 45(4):413–428, 1996.
- [79] S. N. Wood. Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society (B)*, 73(1):3–36, 2011.
- [80] Simon Wood, Fabian Scheipl, and Maintainer Simon Wood. Package ‘gam4’. *Am Stat*, 45(339):0–2, 2017.
- [81] Simon N Wood. Thin plate regression splines. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 65(1):95–114, 2003.
- [82] Yoichi Yokota, Mitsuhiro Aoki, Keisuke Mizuta, Yatsuji Ito, and Naoki Isu. Motion sickness susceptibility associated with visually induced postural instability and cardiac autonomic responses in healthy subjects. *Acta oto-laryngologica*, 125(3):280–285, 2005.